

North Jersey Center for Urologic Care, P.A.

A Division of Urology Group of New Jersey, LLC

16 Pocono Road, Suite 114

Denville, NJ 07834

(Tel) 973-586-3056 (Fax)973-625-0116

www.njcuc.com

Medications to avoid prior to Biopsy or Surgery

You must stop taking the following medications 7-10 days prior to a transrectal ultrasound of the prostate with biopsy. Please do not stop taking other medications prescribed by your doctor. If you have any questions regarding these medications or if you are not sure which medicines or herbal supplements you should stop taking, please call the office at 973-586-3056.

Advil	Cope tablets	Lodine	Rawasa
Alcohol	Cosprin	Magan	Relafen
Aleve	Coumadin	Magsal	Robaxisal
Alka Seltzer	CP-2 Tablets	Major-cin	Rufen
Anacin	Dasin	Majoral	S-A-C Tablets
Anaprox	Daypro	Marnal Caps	Sal-Fayne
Arthralgen	Dinol	Measurin	Salatin
Arthropan	Dipyridamole	Meclomen	Saleto
Arthrotec	Disalcid	Meclofenamate	St. Joseph Cold Tablets
Anaturane	Doan's Pills	Medipren	Sine-off Sinus Medicine
Apap Fort.	Doclin	Mefenamic Acid	Solocol
Argesic	Dolobid	Midol Caplets	Stanback
Artha-G	Duoprin	Micrainin	Supac Caps
Artholate	Duradyne	Mobidin	Synalgos
Aspergum	Durasal	Mobigesic	Tandearil
Aspirbuf	Dynosal	Momentum	Talwin Compound
Aspirin	Easprin	Motrin	Tenstan
Aspiricin	Ecotrin	Nalfon	Ticlid
Asproject	Encaprin	Naprosyn	Tisma
Ascriptin	Efficin	Naproxen	Toradol
Axotal	Emagrin	Neocylate	Trendar
Azulfadine	Empirin	Norgesic Nuprin	Trental
Bayer	Equagesic	Nuprin	Tolectin
BC Tablet and Powder	Excedrin	Orudis	Triaminicin
Buff-A	Fenoprofen	OS-Cal Gesic Tablets	Trigesic
Buffaprin	Fiorinal	Oxalid	Trilisate
Bufferin	Four-Way Cold	Pablate	Ultram
Butazolidin	Garlic	PAC	Uracel
Buffinol	Ginkgo	Panodynes	Vanquish
Buffets II	Gaysal-S	Pepto Bismol	Verin
Buffex	Gemisin	Percodan	Vitamin E Supplement
Buf-Tabs	Goody's HA Powder	Persistin	Voltaren
Cama	Ibuprofen	Plavix	Zorprin
Capron	Indomethacin	Postel	
Celebrex	Indocin	Propoxyphene	
Clinoril	Indo-Lemmon Capsules	Protension	
Congesprin	Lanorinal	Piroxicam (Feldene)	